

Sample exam solutions

Note Title

10/27/2008

①

① $|A| = \frac{1}{2} |4-0| = 2$

$K = 4 - 2 = 2$

↑ ↑
Max |A|

period = π

$\pi = \frac{2\pi}{B}$

$B = 2$

④ phase shift = $\frac{\pi}{4}$

$\frac{\pi}{4} = \frac{-C}{B}$

$\frac{\pi}{4} = \frac{-C}{2}$

$-\frac{\pi}{2} = C$

$y = 2 - 2 \sin\left(2x - \frac{\pi}{2}\right)$

②

① $|A| = \frac{1}{2} |0 - (-6)| = 3$

② $K = 0 - 3 = -3$

③ $A = 3$ not reflected

④ period = $\frac{2\pi}{3}$

$\frac{2\pi}{3} = \frac{2\pi}{B}$

$B = 3$

⑤ p.s. = $\frac{\pi}{6}$

$\frac{\pi}{6} = \frac{-C}{3}$

$-\frac{\pi}{2} = C$

$y = -3 + 3 \sin\left(3x - \frac{\pi}{2}\right)$

③

① $|A| = \frac{1}{2} |0 - (-4)| = 2$

② $K = 0 - 2 = -2$

③ $A = 2$ not reflected

④ period = π

$\pi = \frac{2\pi}{B}$

$B = 2$

⑤ phase shift = $-\frac{\pi}{4}$

$-\frac{\pi}{4} = \frac{-C}{2}$

$\frac{\pi}{2} = C$


$y = -2 + 2 \cos\left(2x + \frac{\pi}{2}\right)$

$$\textcircled{2} \cos^{-1}(\cos(150^\circ)) = 150^\circ \quad 150^\circ \in [0^\circ, 180^\circ]$$

$$\textcircled{b} \tan^{-1}\left(\tan \frac{3\pi}{4}\right) = \tan^{-1}\left(\tan\left(-\frac{\pi}{4}\right)\right) = -\frac{\pi}{4} \quad -\frac{\pi}{4} \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$$


$$\tan\left(\frac{3\pi}{4}\right) = -1$$

$$\tan\left(-\frac{\pi}{4}\right) = -1$$



$$\textcircled{c} \sec\left(\tan^{-1}\left(\frac{3}{4}\right)\right) = \sec \theta = \frac{5}{4}$$

$$\theta = \tan^{-1}\left(\frac{3}{4}\right) \Rightarrow \tan \theta = \frac{3}{4}, \quad 0 \leq \theta < \frac{\pi}{2}$$




$$\textcircled{d} \sec\left(\cos^{-1}\frac{1}{\sqrt{6}}\right) = \frac{1}{\cos\left(\cos^{-1}\left(\frac{1}{\sqrt{6}}\right)\right)} = \frac{1}{\frac{1}{\sqrt{6}}} = \sqrt{6}, \quad \frac{1}{\sqrt{6}} \in [-1, 1]$$

3-7, see book or handouts

9 $\tan(\sin^{-1} 4x)$


$\theta = \sin^{-1} 4x \rightarrow \sin \theta = \frac{4x}{1}$

$$\tan \theta = \frac{4x}{\sqrt{1-16x^2}}$$



Calculator Section

1 $y = 3 + \sin\left(\pi x - \frac{\pi}{2}\right)$



1 $VT = 3$

2 $|A| = 1$

3 $\text{period} = \frac{2\pi}{\pi} = 2, \text{QP} = \frac{1}{2}$

4 one cycle: $0 \leq \pi x - \frac{\pi}{2} \leq 2\pi$
 $\frac{\pi}{2} \leq \pi x \leq \frac{5\pi}{2}$
 $\frac{1}{2} \leq x \leq \frac{5}{2}$

2 $y = -2 + \sec\left(\frac{1}{2}x - \frac{\pi}{3}\right)$

1 $VT = -2$

$|A| = 1$


$\text{period} = \frac{2\pi}{1/2} = 4\pi, \text{QP} = \pi = \frac{3\pi}{3}$

one cycle:

$0 \leq \frac{1}{2}x - \frac{\pi}{3} \leq 2\pi$

$\frac{\pi}{3} \leq \frac{1}{2}x \leq \frac{7\pi}{3}$

$\frac{2\pi}{3} \leq x \leq \frac{14\pi}{3}$



$$(3) y = -1 - \tan\left(\frac{1}{2}x + \frac{\pi}{4}\right)$$

$$(1) vt = -1$$

$$(2) |A| = 1, \text{ reflected}$$


$$(3) \text{ period} = \frac{\pi}{\frac{1}{2}} = 2\pi, \quad OP = \frac{\pi}{2}$$

one cycle:

$$-\frac{\pi}{2} < \frac{1}{2}x + \frac{\pi}{4} < \frac{\pi}{2}$$

$$-\frac{3\pi}{4} < \frac{1}{2}x < \frac{\pi}{4}$$

$$-\frac{3\pi}{2} < x < \frac{\pi}{2}$$




$$(4) (a) \sin 75^\circ = \sin\left(\frac{150^\circ}{2}\right) = \sqrt{\frac{1 - \cos 150^\circ}{2}} = \sqrt{\frac{1 + \frac{\sqrt{3}}{2}}{2}} = \frac{\sqrt{2 + \sqrt{3}}}{2}$$

$$\begin{aligned} \text{or } \sin(45^\circ + 30^\circ) &= \sin 45^\circ \cos 30^\circ + \cos 45^\circ \sin 30^\circ \\ &= \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{3}}{2} + \frac{\sqrt{2}}{2} \cdot \frac{1}{2} \\ &= \frac{\sqrt{6}}{4} + \frac{\sqrt{2}}{4} \end{aligned}$$

$$(b) \tan 105^\circ = \tan\left(\frac{210^\circ}{2}\right) = \frac{1 - \cos 210^\circ}{\sin 210^\circ} = \frac{1 + \frac{\sqrt{3}}{2}}{-\frac{1}{2}} = -\frac{2 + \sqrt{3}}{1} = -2 - \sqrt{3}$$

$$\text{or } \tan(60^\circ + 45^\circ) = \frac{\tan 60^\circ + \tan 45^\circ}{1 - \tan 60^\circ \tan 45^\circ} = \frac{\sqrt{3} + 1}{1 - \sqrt{3}}$$

$$(6) (a) \sin \frac{\theta}{2} = \sqrt{\frac{1 - \cos \theta}{2}} = \sqrt{\frac{1 + \frac{2\sqrt{2}}{3}}{2}} = \frac{\sqrt{3 + 2\sqrt{2}}}{2} \quad \frac{\theta}{2} \in \text{QII}$$




6
b

$$\tan \frac{\theta}{2} = \frac{1 - \cos \theta}{\sin \theta} = \frac{1 - \frac{2\sqrt{2}}{3}}{-\frac{1}{3}} = \frac{3 - 2\sqrt{2}}{-\frac{1}{3}} = -3 + 2\sqrt{2}$$

$$\begin{aligned} \textcircled{7} \quad \cos 2x \sin 8x &= \frac{1}{2} [\sin(2x+8x) - \sin(2x-8x)] \\ &= \frac{1}{2} \sin 10x + \frac{1}{2} \sin 6x, \quad \text{since sine is odd} \end{aligned}$$

$$\begin{aligned} \textcircled{8} \quad \sin 75^\circ - \sin 15^\circ &= 2 \cos \left(\frac{75^\circ + 15^\circ}{2} \right) \sin \left(\frac{75^\circ - 15^\circ}{2} \right) \\ &= 2 \cos \left(\frac{90^\circ}{2} \right) \sin \left(\frac{60^\circ}{2} \right) \\ &= 2 \left(\frac{\sqrt{2}}{2} \right) \cdot \frac{1}{2} \\ &= \frac{\sqrt{2}}{2} \end{aligned}$$

$$\begin{aligned} \textcircled{9} \quad \sin \left(\sec^{-1} \frac{x+1}{3} \right) &= \sin \theta = \frac{\sqrt{x^2+2x-8}}{x+1} \\ \theta &= \sec^{-1} \frac{x+1}{3} \Rightarrow \sec \theta = \frac{x+1}{3} \rightarrow \cos \theta = \frac{3}{x+1} \end{aligned}$$



$$\begin{aligned} \sqrt{(x+1)^2 - 9} &= \sqrt{x^2+2x+1-9} \\ &= \sqrt{x^2+2x-8} \end{aligned}$$

$$\textcircled{10} \quad \frac{\cos t}{1 + \sin t} = \frac{1 - \sin t}{\cos t}$$

LHS


$$\begin{aligned} \frac{\cos t}{1 + \sin t} &= \frac{\cos t (1 - \sin t)}{(1 + \sin t)(1 - \sin t)} = \frac{\cos t (1 - \sin t)}{1 - \sin^2 t} = \frac{\cos t (1 - \sin t)}{\cos^2 t} \\ &= \frac{1 - \sin t}{\cos t} \quad \checkmark \end{aligned}$$

11

$$\textcircled{a} \quad \sqrt{3} + 5 \sin t = 3 \sin t$$

$$2 \sin t = -\sqrt{3}$$


$$\sin t = -\frac{\sqrt{3}}{2}$$




$$\boxed{t = \frac{4\pi}{3} \quad \text{or} \quad t = \frac{5\pi}{3}} \quad 0 < t < 2\pi$$

$$\boxed{t = \frac{4\pi}{3} + 2\pi k \quad \text{or} \quad t = \frac{5\pi}{3} + 2\pi k} \quad \text{all radian solutions}$$

(b) $2 \cos^2 x + \cos x - 1 = 0$
 $(2 \cos x - 1)(\cos x + 1) = 0$
 $\cos x = \frac{1}{2}$ or $\cos x = -1$



$x = \pi/3$ or $x = \frac{5\pi}{3}$



$x = \frac{\pi}{3} + 2\pi k$ or $x = \frac{5\pi}{3} + 2\pi k$


(12)

(a) $\cos 2x - \cos x - 2 = 0$
 $2 \cos^2 x - \cos x - 3 = 0$
 $(2 \cos x - 3)(\cos x + 1) = 0$
 ~~$\cos x = \frac{3}{2}$~~ or $\cos x = -1$
 $\frac{3}{2}$ not in $[-1, 1]$ $x = 180^\circ$

(b) see notes from Section 6.2

(13) $\sin 2x = -\frac{\sqrt{3}}{2}$

$u = 2x$



$\sin u = -\frac{\sqrt{3}}{2}$

$u = 240^\circ + 360^\circ k$ or $u = 300^\circ + 360^\circ k$

$2x = 240^\circ + 360^\circ k$ or $2x = 300^\circ + 360^\circ k$

$x = 120^\circ + 180^\circ k$ or $x = 150^\circ + 180^\circ k$ k any integer

(14)

(a) $x = 2 \cos t$ $y = 2 \sin t$

$\frac{x}{2} = \cos t$ $\frac{y}{2} = \sin t$

$\cos^2 t + \sin^2 t = 1$

$\left(\frac{x}{2}\right)^2 + \left(\frac{y}{2}\right)^2 = 1$

$\frac{x^2}{4} + \frac{y^2}{4} = 1$

$x^2 + y^2 = 4$ circle

(b)

$x = 3 \sin t$ $y = 4 \cos t$

$\frac{x}{3} = \sin t$ $\frac{y}{4} = \cos t$

$\left(\frac{x}{3}\right)^2 + \left(\frac{y}{4}\right)^2 = 1$

$\frac{x^2}{9} + \frac{y^2}{16} = 1$ ellipse

$$\textcircled{c} \quad x = \cos t - 3 \quad y = \sin t + 2$$

$$x + 3 = \cos t \quad y - 2 = \sin t$$

$$(x + 3)^2 + (y - 2)^2 = 1 \quad \text{circle}$$

$$\textcircled{d} \quad x = 3 \cot t \quad y = 3 \csc t$$

$$\frac{x}{3} = \cot t \quad \frac{y}{3} = \csc t$$

$$\cot^2 t + 1 = \csc^2 t$$

$$\csc^2 t - \cot^2 t = 1$$

$$\left(\frac{y}{3}\right)^2 - \left(\frac{x}{3}\right)^2 = 1$$

$$\frac{y^2}{9} - \frac{x^2}{9} = 1 \quad \text{hyperbola}$$